

MINI-REVIEW

Store operated calcium entry in NIH-3T3 cells

Miwako Matsuki-Fukushima^{1,2}, Takuro Tomita², Gary S Bird², and James W Putney, Jr²

¹Nihon University School of Dentistry at Matsudo, Matsudo, Japan ; and ²National Institute of Environmental Health Sciences-NIH, NC, USA

Keywords : store operated calcium entry, NIH-3T3 cells, inhibitors

J. Med. Invest. 56 Suppl. : 381-382, December, 2009

Intracellular Ca²⁺ plays numerous physiological roles in regulating cell functions such as chemotaxis, motility or secretion. Stimulation by neurotransmitters or hormones activates intracellular production of inositol 1, 4, 5-trisphosphate (IP₃) and diacylglycerol (DAG). IP₃ binds to the IP₃ receptor on the Ca²⁺ store (endoplasmic reticulum ; ER) membrane and releases the sequestered Ca²⁺ to the intracellular space. Increased intracellular Ca²⁺ functions as a second messenger for activation of many important downstream effectors. DAG activates protein kinase C (PKC), a key enzyme for diverse physiological functions. Capacitative or store operated calcium entry (SOCE) is a process by which depletion of intracellular Ca²⁺ stores activates a Ca²⁺ influx pathway (1). It is an essential pathway for refilling ER Ca²⁺ stores after releasing store Ca²⁺ and also serves as a direct activator of downstream effectors.

SOCE has been investigated by use of inhibitors of sarcoplasmic-endoplasmic Ca²⁺-ATPase (SERCA), the most common of which is the irreversible inhibitor, thapsigargin. Application of thapsigargin to cells induces sustained SOCE, generally assessed by use of fluorescent Ca²⁺ indicators. Also thapsigargin can be used to study the small, Ca²⁺ selective Ca²⁺ release-activated Ca²⁺ current (*I*_{CRAC}) by the patch clamp technique. Therefore, thapsigargin-induced SOCE is a useful approach to investigate mechanisms of SOCE.

Immortal NIH3T3 cells have been established

from the embryonic NIH Swiss mice fibroblasts. The NIH-3T3 cell is a useful cell line for investigation of transformation, stress fiber formation, or feeder cells of keratinocytes. NIH-3T3 cells have epidermal growth factor (EGF) receptors (2), fibroblast growth factor (FGF) receptors (3), platelet derived growth factor (PDGF) receptors (3), bombesin receptors (3) bradykinin receptors (4) and ATP receptors (5), all of those receptors require calcium influx. Stimulation of those receptors mobilizes stored Ca²⁺ through production of IP₃. However, in a series of papers, it was reported that thapsigargin-induced SOCE in NIH-3T3 cells was not sensitive to several kinds of inhibitors (Table 1) (6-9), indicating that in NIH-3T3 cells, SOCE appears to be independent of PKC, calmodulin, microtubules or actin cytoskeleton.

Table 1 Inhibitors which is insensitive on TG-induced SOCE in NIH-3T3 cells

target	reagent	effect of reagent	reference
protein kinase	staurosporin	inhibition of PKC	8
	PMA	activation of PKC	8
phosphatase	ocadaic acid	inhibition of phosphatase	8
calmodulin	KN62	CAMKII inhibitor	9
	cyclosporin A	calmodulin antagonist	9
cytoskeleton	nocodazole	microtubule inhibitor	6
	cytochalasin D	actin inhibitor	6

Recent discoveries have revealed the major molecular component proteins in SOCE. STromal Interaction Molecule (STIM) 1 and 2 are Ca²⁺ sensing proteins which initiate signaling to open the CRAC

Received for publication October 16, 2009 ; accepted October 23, 2009.

Address correspondence and reprint requests to Miwako Matsuki-Fukushima, Nihon University School of Dentistry at Matsudo, 2-870-1, Sakaecho-Nishi, Matsudo City, Chiba, Japan and Fax : +81-47-360-9327.

channel (10). STIMs have a Ca^{2+} binding site (EF-hand) localized in the N-terminus facing the interior of the Ca^{2+} store. Following depletion of the Ca^{2+} store, the Ca^{2+} dissociates from the EF-hand and STIMs are activated. Activated STIMs then interact with the CRAC channel and to open the channel pore. After the discovery of STIMs, Orai proteins were found to be essential partners of STIM proteins (10). Orai1, 2 and 3 are believed to function as pore-forming subunits of the store-operated channels. The Orai1 channel is blocked by low concentrations of Gd^{3+} ($\sim 1 \mu\text{M}$) or by of 2-aminoethoxydiphenyl borate (2-APB) (11). In the future, molecular manipulations of Orai and STIM expression, as well as use of Gd^{3+} or 2-APB will be useful tools to investigate the mechanism and regulation of SOCE in NIH-3T3 cells.

ACKNOWLEDGEMENTS

Work discussed in this review originating in the authors' laboratory was supported by the Intramural Program, National Institutes of Health.

REFERENCES

1. Putney JW Jr : A model for receptor-regulated calcium entry. *Cell Calcium* 7 : 1-12, 1986
2. Moolenaar WH, Bierman AJ, Tilly BC, Verlaan I, Defize LH, Honegger AM, Ullrich A, Schlessinger J : A point mutation at the ATP-binding site of the EGF-receptor abolishes signal transduction. *EMBO J* 7 : 707-10, 1988
3. Kojima I, Nishimoto I, Iiri T, Ogata E, Rosenfeld R : Evidence that type II insulin-like growth factor receptor is coupled to calcium gating system. *Biochem Biophys Res Commun* 154 : 9-19, 1988
4. Magni M, Meldolesi J, Pandiella A : Ionic events induced by epidermal growth factor. Evidence that hyperpolarization and stimulated cation influx play a role in the stimulation of cell growth. *J Biol Chem* 266 : 6329-35, 1991
5. Giovannardi S, Racca C, Bertollini L, Sturani E, Peres A : P2Y purinoceptors in normal NIH 3T3 and in NIH 3T3 overexpressing c-ras. *Exp Cell Res* 202 : 398-404, 1992
6. Ribeiro CM, Reece J, Putney JW Jr : Role of the cytoskeleton in calcium signaling in NIH 3T3 cells. An intact cytoskeleton is required for agonist-induced $[\text{Ca}^{2+}]_i$ signaling, but not for capacitative calcium entry. *J Biol Chem* 272 : 26555-61, 1997
7. Lobaugh LA, Eisfelder B, Gibson K, Johnson GL, Putney JW Jr : Constitutive activation of a phosphoinositidase C-linked G protein in murine fibroblasts decreases agonist-stimulated Ca^{2+} mobilization. *Mol Pharmacol* 50 : 493-500, 1996
8. Ribeiro CM, Putney JW Jr : Differential effects of protein kinase C activation on calcium storage and capacitative calcium entry in NIH 3T3 cells. *J Biol Chem* 271 : 21522-8, 1996
9. Louzao MC, Ribeiro CM, Bird GS, Putney JW Jr : Cell type-specific modes of feedback regulation of capacitative calcium entry. *J Biol Chem* 271 : 14807-13, 1996
10. Putney JW Jr : New molecular players in capacitative Ca^{2+} entry. *J Cell Sci* 120 : 1959-65, 2007
11. DeHaven WI, Smyth JT, Boyles RR, Bird GS, Putney JW Jr : Complex actions of 2-aminoethoxydiphenyl borate on store-operated calcium entry. *J Biol Chem* 283 : 19265-73, 2008