

## ORIGINAL

# Y chromosome haplogroups are associated with birth size in Japanese men

Youichi Sato

*Department of Pharmaceutical Information Science, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan*

**Abstract:** The Y chromosome is classified into haplogroups (A-T) based on a combination of several DNA polymorphisms. Japanese men are mainly classified into haplogroups C, D, and O, which have been further subdivided. The distribution of Y-chromosome haplogroups varies by ethnicity. The phylogenetic age, origin, and migration also differ. I hypothesized that Y chromosome haplogroups may be associated with height and/or weight at birth. An association analysis of height and weight at birth with Y chromosome haplogroups was performed in 288 Japanese men. Men belonging to haplogroup O1b2 were significantly associated with short stature at birth (beta = -1.88, standard error (SE) = 0.55,  $P = 0.00076$ ), and those belonging to D1a2a-12f2b were significantly associated with increased birth weight (beta = 174, SE = 64,  $P = 0.0069$ ). Y chromosome haplogroups are associated with physical birth characteristics in modern Japanese men. *J. Med. Invest.* 71:129-133, February, 2024

**Keywords:** birth height, birth weight, Japanese population, Y chromosome haplogroup

## INTRODUCTION

The Y chromosome does not undergo genetic recombination and is passed on to the son as an intact chromosome. Occasional mutations occur, and combinations of DNA polymorphisms are classified into haplogroups (A-T) (1). Haplogroups are characterized by ethnicity and region, with Japanese men primarily classified into haplogroups C, D, or O (2). Haplogroups C and D were reported to have expanded into Japan approximately 12,000 to 20,000 years ago, and haplogroup O is subdivided into O1b and O2a, of which O1b was reported to have expanded 4,000 years ago (3, 4). Therefore, I hypothesized that there may be a relationship between the Y chromosome haplogroup and the physical characteristics of Japanese men.

Recently, a genome-wide association study (GWAS) of 5.4 million individuals reported 12,111 independent single nucleotide polymorphisms (SNPs) significantly associated with height, representing nearly all of the SNP-based common heritability (5). Thus, human height is largely a function of genetic background. In addition, many GWAS of body weight have been conducted, and more than 1,100 loci associated with obesity have been reported (6).

The physical characteristics of adults vary greatly depending on environmental factors such as age and lifestyle. However, compared with adults, the physical characteristics of children at birth are thought to be less influenced by the environment. In the present study, I examined the relationship between Y chromosome haplogroups and birth size (height and weight) in Japanese men.

## MATERIALS AND METHODS

This study was approved by the Ethics Committees of the University of Tokushima and St. Marianna Medical University.

### *Samples from Japanese men and genotype of Y chromosome haplogroup*

The subjects have been described previously (2, 7). Briefly, participants were recruited from the partners of pregnant women attending obstetric clinics in Sapporo, Osaka, Kanazawa and Fukuoka cities in Japan from 1999 to 2002. Multiple cohorts were analyzed to increase the sample size. Of the 792 participants, 288 Japanese men who self-reported birth weight and height were included in this study. Supermales or other syndromes that affect height were not tested. The Y chromosomes of all subjects were defined as 15 haplogroups according to the Y Chromosome Consortium (YCC) nomenclature (1) using 24 Y chromosome markers (YAP, M174, M15, M55, 12f2b, RSP4Y, M8, M217, M213, M9, M214, LLY22g, M175, MSY2.2, M119, P31, M95, SRY465, 47z, M122, M324, JST021354, M134, and JST002611), as previously described (2). The primer sets, PCR amplification conditions, and restriction enzymes are also described previously (2). The haplogroups were renamed according to the International Society of Genetic Genealogy (ISOGG) phylogenetic tree on April 19, 2019 (<https://isogg.org/tree/>).

### *Statistical analysis*

Associations between Y chromosome haplogroups with frequencies > 0.05 and birth weight and height were assessed using linear regression analysis. No adjustment for covariates was made, because the only variables were height and weight. Statistical analyses were performed using R version 4.0.3 (The R Project for Statistical Computing; <http://www.r-project.org>), and statistical significance was set at  $P < 0.025$  (0.05/2 test: birth height and weight) to account for multiple testing.

Received for publication October 29, 2023; accepted December 27, 2023.

Address correspondence and reprint requests to Youichi Sato, Ph.D., Department of Pharmaceutical Information Science, Tokushima University Graduate School of Biomedical Sciences, 1-78-1 Sho-machi, Tokushima, Tokushima 770-8505, Japan and Fax: +81-88-633-7253. E-mail: youichi.sato@tokushima-u.ac.jp

## RESULTS

*Y chromosome haplogroups and birth height and weight in Japanese men*

The mean birth height and weight of the 288 subjects were  $50.1 \pm 2.5$  cm and  $3180.5 \pm 399.1$  g, respectively. To investigate the association between Y chromosome haplogroups and birth height and weight, I typed Y chromosome haplogroups using 24 Y chromosome markers. Participants belonged to 15 haplogroups and their haplogroup distributions are shown in Table 1. The haplogroup O1b2a1a1 was the most dominant, and C1a1, C2, D1a2a-M55, D1a2a-12f2b, O1b2, O2a2b1, and O2a1b had

frequencies > 5% among the participants. The frequency distributions of the subjects was similar to those of other cohorts (8-11). The means and standard deviations of birth weight and height for each haplotype are also shown in Table 1.

I then analyzed the associations between these eight Y chromosome haplogroups and birth weight and height using linear regression analysis. I found that men belonging to haplogroup O1b2 were significantly associated with shorter stature at birth (beta = -1.88, standard error (SE) = 0.55,  $P = 0.00076$ ) than other haplogroups, and men belonging to D1a2a-12f2b were significantly associated with higher birth weight (beta = 174, SE = 64,  $P = 0.0069$ ) than other haplogroups (Table 2). Birth

**Table 1.** Numbers and percent frequencies of Y chromosome haplogroups and means and standard deviations of birth weight and height for each haplotype in 288 Japanese men

Y chromosome haplogroup	Numbers (%)	Birth height (cm)		Birth weight (g)	
		Mean	SD	Mean	SD
C1a1	17 (5.9)	50.2	2.2	3082.1	530.5
C2	22 (7.6)	49.9	1.7	3233.0	311.5
D1a2a-M55	37 (12.8)	49.5	2.8	3079.1	449.0
D1a2a-12f2b	45 (15.6)	50.4	2.0	3327.4	412.2
F*	2 (0.7)	48.5	2.1	2601.5	252.4
K*	12 (4.2)	51.1	2.3	3146.7	368.4
N1	4 (1.4)	49.3	0.5	3005.0	310.4
O1a	6 (2.1)	50.6	2.6	3276.7	461.2
O1b1a1a	3 (1.0)	49.5	2.5	3063.3	77.7
O1b2	21 (7.3)	48.3	3.4	3081.3	374.9
O1b2a1a1	63 (21.9)	50.4	2.7	3188.6	324.3
O2	7 (2.4)	49.5	2.1	2932.1	428.6
O2a2	8 (2.8)	49.7	3.1	3152.1	287.6
O2a2b1	24 (8.3)	50.7	2.3	3254.6	420.5
O2a1b	17 (5.9)	50.5	1.4	3265.9	436.5

**Table 2.** Association analysis between Y chromosome haplogroup and height and weight at birth in 288 Japanese men

Y chromosome haplogroup	Birth height			Birth weight		
	Beta	(SE)	<i>P</i> -value	Beta	(SE)	<i>P</i> -value
C1a1	0.11	(0.62)	0.86	-105	(100)	0.30
C2	-0.22	(0.55)	0.69	57	(89)	0.52
D1a2a-M55	-0.71	(0.44)	0.10	-116	(70)	0.098
<b>D1a2a-12f2b</b>	0.44	(0.40)	0.27	<b>174</b>	<b>(64)</b>	<b>0.0069</b>
F*	-1.53	(1.76)	0.38	-583	(281)	0.039
K*	1.10	(0.73)	0.13	-35	(118)	0.77
N1	-0.84	(1.25)	0.50	-178	(201)	0.38
O1a	0.52	(1.02)	0.61	98	(165)	0.55
O1b1a1a	-0.58	(1.44)	0.69	-118	(232)	0.61
<b>O1b2</b>	<b>-1.88</b>	<b>(0.55)</b>	<b>0.00076</b>	-107	(90)	0.24
O1b2a1a1	0.46	(0.35)	0.19	10	(57)	0.86
O2	-0.56	(0.95)	0.56	-255	(152)	0.096
O2a2	-0.40	(0.89)	0.66	-29	(143)	0.84
O2a2b1	-0.65	(0.53)	0.22	81	(85)	0.34
O2a1b	0.50	(0.62)	0.42	91	(100)	0.36

Data are shown as estimated linear regression statistic beta, standard error (SE), and *P*-value. Numbers in bold indicate significance ( $P < 0.025$ ).

weight is known to have decreased over the last 30 years (12). To confirm that age may be a factor, I analyzed age as a covariate. The association between men belonging to D1a2a-12f2b and higher birth weight remained significant after adjusting for age (beta = 179, SE = 64,  $P = 0.0051$ ).

## DISCUSSION

In this study, compared with men belonging to the Y chromosome haplogroups, those belonging to the O1b2 haplogroup were associated with shorter birth height, and those belonging to haplogroup D1a2a-12f2b were associated with higher birth weight in the Japanese population. Haplogroup O1b2 expanded approximately 4,000 years ago (3, 4) and is frequently found in Korea outside of Japan (13). To date, the phenotypic manifestation of O1b2 has been unclear; this study is the first to show that O1b2 is associated with short stature at birth. The O1b2 marker SRY465 is an SNP with a synonymous substitution in the sex-determining region Y (*SRY*) gene (14). *SRY* is required for male determination; however, it is unclear how it relates to height at birth. A gene on the Y chromosome linked to SRY465 may influence birth height; however, further studies are required to clarify this phenomenon.

D1a2a-12f2b, which has been shown to be associated with higher birth weight, is a haplogroup found only in the Japanese population. Previously, we found that men belonging to haplogroup D1a2a-12f2b had higher levels of luteinizing hormone (LH) (15) and lower semen quality (16). Furthermore, the frequency of haplogroup D1a2a-12f2b was found to be significantly higher in patients with azoospermia than in controls (17). LH is a hormone secreted by the pituitary gland in the brain that contributes to sperm production by inducing testosterone biosynthesis and secretion in testicular Leydig cells (18, 19). These results indicate that men belonging to haplogroup D1a2a-12f2b have low spermatogenic potential. Several studies have reported the association between birth weight and infertility in men. Francois *et al.* reported that low birth weight correlated with unexplained subfertility (20). Kahn *et al.* reported that birth weight percentile during gestation was positively correlated with square root sperm concentration (21). Therefore, low birth weight appears to be associated with infertility and poor sperm quality. However, this was in contrast to our finding that men belonging to haplogroup D2a1 were associated with lower sperm quality and higher weight at birth. This discrepancy may be due to differences between the epidemiological and genetic association studies.

A limitation of this study is that it relies on self-reports of birth weight and length, and it is unclear how accurate these are. Data collection on important factors that may influence variables such as length of pregnancy and maternal health is also lacking. In addition, only 288 of the original sample of 792 adult males responded to the survey. The use of 288 samples in 15 haplogroups is considered a reasonable number compared to a sample size of over 100,000 SNPs, as in a GWAS. However, the sample size also needs to be increased and will be addressed in future studies.

## CONCLUSIONS

Although several limitations need to be overcome, this study provides the first evidence of a relationship between Y chromosome haplogroups and physical characteristics in Japanese men.

## CONFLICT OF INTEREST STATEMENT

The author declares no conflict of interest.

## ACKNOWLEDGEMENTS

Yutaka Nakahori, Toshikatsu Shinka, Eitetsue Koh, Jiro Kanaya, Mikio Namiki, Kiyomi Matsumiya, Akira Tsujimura, Kiyoshi Komatsu, Naoki Itoh, and Jiro Eguchi for collecting blood samples from participants. This study was supported by the Grants-in-Aid for Scientific Research on Innovative Areas "Yaponesian Genome" (19H05346) from the Japan Society for the Promotion of Science. I would like to thank Editage (www.editage.com) for the English language editing.

## REFERENCES

- Karafet TM, Mendez FL, Meilerman MB, Underhill PA, Zegura SL, Hammer MF: New binary polymorphisms reshape and increase resolution of the human Y chromosomal haplogroup tree. *Genome Res* 18 : 830-838, 2008
- Sato Y, Shinka T, Ewis AA, Yamauchi A, Iwamoto T, Nakahori Y: Overview of genetic variation in the Y chromosome of modern Japanese males. *Anthropological Science* 122 : 131-136, 2014
- Hammer MF, Karafet TM, Park H, Omoto K, Harihara S, Stoneking M, Horai S: Dual origins of the Japanese: common ground for hunter-gatherer and farmer Y chromosomes. *J Hum Genet* 51 : 47-58, 2006
- Hammer MF, Horai S: Y chromosomal DNA variation and the peopling of Japan. *Am J Hum Genet* 56 : 951-962, 1995
- Yengo L, Vedantam S, Marouli E, Sidorenko J, Bartell E, Sakaue S, Graff M, Eliassen AU, Jiang Y, Raghavan S, Miao J, Arias JD, Graham SE, Mukamel RE, Spracklen CN, Yin X, Chen SH, Ferreira T, Highland HH, Ji Y, Karaderi T, Lin K, Lüll K, Malden DE, Medina-Gomez C, Machado M, Moore A, Rieger S, Sim X, Vrieze S, Ahluwalia TS, Akiyama M, Allison MA, Alvarez M, Andersen MK, Ani A, Appadurai V, Arbeeve L, Bhaskar S, Bielak LF, Bollepalli S, Bonnycastle LL, Bork-Jensen J, Bradfield JP, Bradford Y, Braund PS, Brody JA, Burgdorf KS, Cade BE, Cai H, Cai Q, Campbell A, Cañadas-Garre M, Catamo E, Chai JF, Chai X, Chang LC, Chang YC, Chen CH, Chesani A, Choi SH, Chung RH, Cocca M, Concas MP, Couture C, Cuellar-Partida G, Danning R, Daw EW, Degenhard F, Delgado GE, Delitala A, Demirkan A, Deng X, Devineni P, Dietl A, Dimitriou M, Dimitrov L, Dorajoo R, Ekici AB, Engmann JE, Fairhurst-Hunter Z, Farmaki AE, Faul JD, Fernandez-Lopez JC, Forer L, Francescatti M, Freitag-Wolf S, Fuchsberger C, Galesloot TE, Gao Y, Gao Z, Geller F, Giannakopoulou O, Giulianini F, Gjesing AP, Goel A, Gordon SD, Gorski M, Grove J, Guo X, Gustafsson S, Haessler J, Hansen TF, Havulinna AS, Haworth SJ, He J, Heard-Costa N, Hebbar P, Hindy G, Ho YA, Hofer E, Holliday E, Horn K, Hornsby WE, Hottenga JJ, Huang H, Huang J, Huerta-Chagoya A, Huffman JE, Hung YJ, Huo S, Hwang MY, Iha H, Ikeda DD, Isono M, Jackson AU, Jäger S, Jansen IE, Johansson I, Jonas JB, Jonsson A, Jørgensen T, Kalafati IP, Kanai M, Kanoni S, Kärhus LL, Kasturiratne A, Katsuya T, Kawaguchi T, Kember RL, Kentistou KA, Kim HN, Kim YJ, Kleber ME, Knol MJ, Kurbasic A, Lauzon M, Le P, Lea R, Lee JY, Leonard HL, Li SA, Li X, Li X, Liang J, Lin H, Lin SY, Liu J, Liu X, Lo KS, Long J, Lores-Motta L, Luan J, Lyssenko V,

- Lytyikäinen LP, Mahajan A, Mamakou V, Mangino M, Manichaikul A, Marten J, Mattheisen M, Mavarani L, McDaid AF, Meidtner K, Melendez TL, Mercader JM, Milaneschi Y, Miller JE, Millwood IY, Mishra PP, Mitchell RE, Møllehave LT, Morgan A, Mucha S, Munz M, Nakatochi M, Nelson CP, Nethander M, Nho CW, Nielsen AA, Nolte IM, Nongmaithem SS, Noordam R, Ntalla I, Nutile T, Pandit A, Christofidou P, Pärna K, Pauper M, Petersen ERB, Petersen LV, Pitkänen N, Polašek O, Poveda A, Preuss MH, Pyarajan S, Raffield LM, Rakugi H, Ramirez J, Rasheed A, Raven D, Rayner NW, Riveros C, Rohde R, Ruggiero D, Ruotsalainen SE, Ryan KA, Sabater-Lleal M, Saxena R, Scholz M, Sendamarai A, Shen B, Shi J, Shin JH, Sidore C, Sitlani CM, Sliker RC, Smit RAJ, Smith AV, Smith JA, Smyth LJ, Southam L, Steinthorsdottir V, Sun L, Takeuchi F, Tallapragada DSP, Taylor KD, Tayo BO, Tcheandjieu C, Terzikhan N, Tesolin P, Teumer A, Theusch E, Thompson DJ, Thorleifsson G, Timmers PRHJ, Trompet S, Turman C, Vaccargiu S, van der Laan SW, van der Most PJ, van Klinken JB, van Setten J, Verma SS, Verweij N, Veturi Y, Wang CA, Wang C, Wang L, Wang Z, Warren HR, Bin Wei W, Wickremasinghe AR, Wielscher M, Wiggins KL, Winsvold BS, Wong A, Wu Y, Wuttke M, Xia R, Xie T, Yamamoto K, Yang J, Yao J, Young H, Yousri NA, Yu L, Zeng L, Zhang W, Zhang X, Zhao JH, Zhao W, Zhou W, Zimmermann ME, Zoledziwska M, Adair LS, Adams HHH, Aguilar-Salinas CA, Al-Mulla F, Arnett DK, Asselbergs FW, Åsvold BO, Attia J, Banas B, Bandinelli S, Bennett DA, Bergler T, Bharadwaj D, Biino G, Bisgaard H, Boerwinkle E, Böger CA, Bønnelykke K, Boomsma DI, Børghlum AD, Borja JB, Bouchard C, Bowden DW, Brandslund I, Brumpton B, Buring JE, Caulfield MJ, Chambers JC, Chandak GR, Chanock SJ, Chaturvedi N, Chen YI, Chen Z, Cheng CY, Christophersen IE, Ciullo M, Cole JW, Collins FS, Cooper RS, Cruz M, Cucca F, Cupples LA, Cutler MJ, Damrauer SM, Dantoft TM, de Borst GJ, de Groot LCPGM, De Jager PL, de Kleijn DPV, Janaka de Silva H, Dedoussis GV, den Hollander AI, Du S, Easton DF, Elders PJM, Eliassen AH, Ellinor PT, Elmståhl S, Erdmann J, Evans MK, Fatkin D, Feenstra B, Feitosa MF, Ferrucci L, Ford I, Fornage M, Franke A, Franks PW, Freedman BI, Gasparini P, Gieger C, Girotto G, Goddard ME, Golightly YM, Gonzalez-Villalpando C, Gordon-Larsen P, Grallert H, Grant SFA, Grarup N, Griffiths L, Gudnason V, Haiman C, Hakonarson H, Hansen T, Hartman CA, Hattersley AT, Hayward C, Heckbert SR, Heng CK, Hengstenberg C, Hewitt AW, Hishigaki H, Hoyng CB, Huang PL, Huang W, Hunt SC, Hveem K, Hyppönen E, Iacono WG, Ichihara S, Ikram MA, Isasi CR, Jackson RD, Jarvelin MR, Jin ZB, Jöckel KH, Joshi PK, Jousilahti P, Jukema JW, Kähönen M, Kamatani Y, Kang KD, Kaprio J, Kardia SLR, Karpe F, Kato N, Kee F, Kessler T, Khera AV, Khor CC, Kiemeny LALM, Kim BJ, Kim EK, Kim HL, Kirchhof P, Kivimaki M, Koh WP, Koistinen HA, Kolovou GD, Kooner JS, Kooperberg C, Köttgen A, Kovacs P, Kraaijeveld A, Kraft P, Krauss RM, Kumari M, Kutalik Z, Laakso M, Lange LA, Langenberg C, Launer LJ, Le Marchand L, Lee H, Lee NR, Lehtimäki T, Li H, Li L, Lieb W, Lin X, Lind L, Linneberg A, Liu CT, Liu J, Loeffler M, London B, Lubitz SA, Lye SJ, Mackey DA, Mägi R, Magnusson PKE, Marcus GM, Vidal PM, Martin NG, März W, Matsuda F, McGarrah RW, McGue M, McKnight AJ, Medland SE, Mellström D, Metspalu A, Mitchell BD, Mitchell P, Mook-Kanamori DO, Morris AD, Mucci LA, Munroe PB, Nalls MA, Nazarian S, Nelson AE, Neville MJ, Newton-Cheh C, Nielsen CS, Nöthen MM, Ohlsson C, Oldehinkel AJ, Orozco L, Pahlkala K, Pajukanta P, Palmer CNA, Parra EJ, Pattaro C, Pedersen O, Pennell CE, Penninx BWJH, Perusse L, Peters A, Peyser PA, Porteous DJ, Posthuma D, Power C, Pramstaller PP, Province MA, Qi Q, Qu J, Rader DJ, Raitakari OT, Ralhan S, Rallidis LS, Rao DC, Redline S, Reilly DF, Reiner AP, Rhee SY, Ridker PM, Rienstra M, Ripatti S, Ritchie MD, Roden DM, Rosendaal FR, Rotter JI, Rudan I, Rutter F, Sabanayagam C, Saleheen D, Salomaa V, Samani NJ, Sanghera DK, Sattar N, Schmidt B, Schmidt H, Schmidt R, Schulze MB, Schunkert H, Scott LJ, Scott RJ, Sever P, Shiroma EJ, Shoemaker MB, Shu XO, Simonsick EM, Sims M, Singh JR, Singleton AB, Sinner MF, Smith JG, Snieder H, Spector TD, Stampfer MJ, Stark KJ, Strachan DP, 't Hart LM, Tabara Y, Tang H, Tardif JC, Thanaraj TA, Timpson NJ, Tönjes A, Tremblay A, Tuomi T, Tuomilehto J, Tusié-Luna MT, Uitterlinden AG, van Dam RM, van der Harst P, Van der Velde N, van Duijn CM, van Schoor NM, Vitart V, Völker U, Vollenweider P, Völzke H, Wacher-Rodarte NH, Walker M, Wang YX, Wareham NJ, Watanabe RM, Watkins H, Weir DR, Werge TM, Widen E, Wilkens LR, Willemsen G, Willett WC, Wilson JF, Wong TY, Woo JT, Wright AF, Wu JY, Xu H, Yajnik CS, Yokota M, Yuan JM, Zeggini E, Zemel BS, Zheng W, Zhu X, Zmuda JM, Zonderman AB, Zwart JA ; 23andMe Research Team ; VA Million Veteran Program ; DiscovEHR (DiscovEHR and MyCode Community Health Initiative) ; eMERGE (Electronic Medical Records and Genomics Network) ; Lifelines Cohort Study ; PRACTICAL Consortium ; Understanding Society Scientific Group ; Chasman DI, Cho YS, Heid IM, McCarthy MI, Ng MCY, O'Donnell CJ, Rivadeneira F, Thorsteinsdottir U, Sun YV, Tai ES, Boehnke M, Deloukas P, Justice AE, Lindgren CM, Loos RJF, Mohlke KL, North KE, Stefansson K, Walters RG, Winkler TW, Young KL, Loh PR, Yang J, Esko T, Assimes TL, Auton A, Abecasis GR, Willer CJ, Locke AE, Berndt SI, Lettre G, Frayling TM, Okada Y, Wood AR, Visscher PM, Hirschhorn JN : A saturated map of common genetic variants associated with human height. *Nature* 610 : 704-712, 2022
6. Loos RJF, Yeo GSH : The genetics of obesity : from discovery to biology. *Nat Rev Genet* 23 : 120-133, 2022
  7. Iwamoto T, Nozawa S, Yoshiike M, Namiki M, Koh E, Kanaya J, Okuyama A, Matsumiya K, Tsujimura A, Komatsu K, Tsukamoto T, Itoh N, Mieno MN, Vierula M, Toppari J, Skakkebaek NE, Jorgensen N : Semen quality of fertile Japanese men : a cross-sectional population-based study of 792 men. *BMJ Open* 3 : e002223, 2013
  8. Tajima A, Hayami M, Tokunaga K, Juji T, Matsuo M, Marzuki S, Omoto K, Horai S : Genetic origins of the Ainu inferred from combined DNA analyses of maternal and paternal lineages. *J Hum Genet* 49 : 187-193, 2004
  9. Hammer MF, Karafet TM, Park H, Omoto K, Harihara S, Stoneking M, Horai S : Dual origins of the Japanese : common ground for hunter-gatherer and farmer Y chromosomes. *J Hum Genet* 51 : 47-58, 2006
  10. Naitoh S, Kasahara-Nonaka I, Minaguchi K, Nambiar P : Assignment of Y-chromosomal SNPs found in Japanese population to Y-chromosomal haplogroup tree. *J Hum Genet* 58 : 195-201, 2013
  11. Harayama Y, Kamei S, Sato N, Hayashi T, Shiozaki T, Ota M, Asamura H : Analysis of Y chromosome haplogroups in Japanese population using short amplicons and its application in forensic analysis. *Leg Med (Tokyo)* 16 : 20-25, 2014
  12. Kato N, Sauvaget C, Yoshida H, Yokoyama T, Yoshiike N : Factors associated with birthweight decline in Japan (1980-2004). *BMC Pregnancy Childbirth* 21 : 337, 2021

13. Kim SH, Kim KC, Shin DJ, Jin HJ, Kwak KD, Han MS, Song JM, Kim W, Kim W : High frequencies of Y-chromosome haplogroup O2b-SRY465 lineages in Korea : a genetic perspective on the peopling of Korea. *Investig Genet* 2 : 10, 2011
14. Shinka T, Tomita K, Toda T, Kotliarova SE, Lee J, Kuroki Y, Jin DK, Tokunaga K, Nakamura H, Nakahori Y : Genetic variations on the Y chromosome in the Japanese population and implications for modern human Y chromosome lineage. *J Hum Genet* 44 : 240-245, 1999
15. Sato Y, Shinka T, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Yamauchi A, Iwamoto T, Nakahori Y : Y chromosome haplogroup D2a1 is significantly associated with high levels of luteinizing hormone in Japanese men. *Andrology* 3 : 520-525, 2015
16. Sato Y, Iwamoto T, Shinka T, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Yamauchi A, Nakahori Y : Y chromosome gr/gr subdeletion is associated with lower semen quality in young men from the general Japanese population but not in fertile Japanese Men. *Biol Reprod* 90 : 116, 2014
17. Sato Y, Shinka T, Iwamoto T, Yamauchi A, Nakahori Y : Y chromosome haplogroup d2\* lineage is associated with azoospermia in Japanese males. *Biol Reprod* 88 : 107, 2013
18. Belchetz PE, Plant TM, Nakai Y, Keogh EJ, Knobil E : Hypophysial responses to continuous and intermittent delivery of hypothalamic gonadotropin-releasing hormone. *Science* 202 : 631-633, 1978
19. Smith MA, Vale WW : Desensitization to gonadotropin-releasing hormone observed in superfused pituitary cells on Cytodex beads. *Endocrinology* 108 : 752-759, 1981
20. Francois I, de Zegher F, Spiessens C, D'Hooghe T, Vanderschueren D : Low birth weight and subsequent male subfertility. *Pediatr Res* 42 : 899-901, 1997
21. Kahn LG, Widen EM, Janevic T, Straka N, Liu X, Cirillo PM, Cohn BA, Buck Louis GM, Factor-Litvak P : The Relation of Birth Weight and Adiposity Across the Life Course to Semen Quality in Middle Age. *Epidemiology* 30 Suppl 2(Suppl 2) : S17-S27, 2019